Pregunta: Qué es esto?




Fe2+
Proximal His
Distal His
Side view
Myoglobin with heme
Heme 90° rotation


A Dash of Science.com
Meat Color
myoglobin
myoglobin is not holding oxygen
Fe+2
globin
oxymyoglobin
myoglobin is holding oxygen
O2
Fe+2
globin
metmyoglobin
myoglobin is holding water
H2O
Fe+3
globin

Química de Coordinación

UNAM Mayo 24 y 26, 2022

peter.kroneck@uni-konstanz.de
https://www.researchgate.net/profile/Peter-Kroneck

Iones metálicos en sistemas vivos
Metaloenzimas y Metaloproteínas

Chlorophyll a



Heme b



Trayendo química inorgánica a vida..

Biochemistry

Biogeochemistry

Physiology

Molecular Biology

Biophysics

Química de Coordinación Bioinorgánica

Microbiology

Environmental Chemistry

Spectroscopy

Structural Biology

Catalysis

Physical Chemistry

Medical Chemistry/Toxicology

Metales y Vida:
la Química de Coordinación de la Naturaleza

“El uso de metales para tratar dolencias humanas se remonta al menos al quinto siglo a. de J.C., y el estudio y la imitación de metales en la biología son un sujeto vibrante hoy”

Stephen Lippard, J Am Chem Soc (2010), 132, 141689-14693





Cisplatin-DNA adduct bound to HMGB1. Cisplatin shown as red and white spheres; DNA is shown in blue; HMBG1 shown as grey cartoon with intercalated phenylalanine shown as yellow spheres. Image credits: Michael S. McCormick.



B. Rosenberg et al., (1965) Nature, 205, 698 - 699

Metales en Medicina – Aplicaciones

"Uno de los desafíos de diseñar medicinas basadas en el metal es equilibrar la toxicidad potencial de una formulación activa con el impacto positivo sustancial de estos recursos terapéuticos y diagnósticos cada vez más comunes"

K.H. Thompson, C. Orvig (2003) Science <u>300</u>, 936-939



Los elementos/metales de vida

www.webelements.com





Mo: 9 mg

METAL DE TRANSICION

Bibliografia Basica y Complementaria

J. J. R. Fraústo da Silva, R. J. P. Williams, 2001
The biological chemistry of the elements, Oxford University Press

G.A. Lawrance, 2010
Introduction to Coordination Chemistry, Wiley

R. R. Crichton, 2008, 2012 y 2019
Biological Inorganic Chemistry, Elsevier

Chemical Reviews, 1996; 2004
Special Issues on Bioinorganic Enzymology, *96*, 2237; *104*, 347

E. R. Featherston, J. A. Cotruvo, 2021 (Review)
The biochemistry of lanthanide acquisition, trafficking, and utilization
Biochim Biophys Acta, Molecular Cell Research *1868*, 118864
https://doi.org/10.1016/j.bbamcr.2020.118864

Sitios Web Importantes

https://www.ebi.ac.uk/pdbe/

comprehensive database of all published protein structures

http://www.brenda-enzymes.org/

comprehensive enzyme database including information on metal requirements

http://www.webelements.com

periodic table of the elements including useful information on each element

http://www.rsc.org/chemsoc/visualelements/pages/alchemist/alchemy.html

PDB Database

- **PDB = Protein Data Bank https://www.rcsb.org/; type in the search field the PDB number**

- **1A70 (for Ferredoxin)**
- **3SBP (for Nitrous Oxide Reductase)**


(a)
AlaX-M (**Zn**, **Ni**)
PDB:2E1B
Zinc finger (**Zn**, **Ni**)
PDB: 1ZAA
Zinc hook (**Zn**, **Hg**)
Rad50, PDB: 1L8D
Gal4 (**Zn**, **Cd**)
PDB:1D66
Cu/**Zn** superoxide
dismutase; PDB: 3F7K
(b)
2**Fe**-2S cluster
(SoxR, PDB: 2ZHH)
4**Fe**-4S cluster
(XPD, PDB: 2CRV)
(c)
Vitamin B12 (**Cu**)
Cobalamin, PDB: 2BB6
Fe**Mo**Co (**V**)
PDB: 1H1L
Heme (**Fe**)
Cytochrome, PDB: 3OFT

Por qué investigan metales en la biología?

- **Hay apenas cualquier proceso importante en la naturaleza que no depende de un <u>ión</u> metálico; ~ 1/3 de las proteínas del genoma humano dependen de <u>iones</u> metálicos**
- **Dos ejemplos importantes:**
- **Catálisis ácida y baja**
- **Química de redox – Transferencia de protones y electrones (conservación de energía)**

Transferencia de protones y electrones
Conservación de energía



$$\Delta G^{o\prime} = -nF\Delta E^{o\prime}$$

Respiración de Mitochondrial & Síntesis de Centro de Reacción Fotosintética de ATP – transferencia de electrón/protón Conectada – fuerza de Protonmotive









Objetivo: De Estructura 3D & Electrónica muy resuelta para Funcionar a Mecanismo



Por qué (transición) iones de metal ?

- **<u>Positively Charged</u>**
  - **Lewis Acids**
  - **Stabilization of Anions**
- **<u>Loosely Bound Electrons</u>**
  - **Redox Active**
  - **Multiple Redox States**
  - **Easily tunable Redox Potential**
- **<u>Redox/Acid Base Chemistry</u>**

- **<u>Open Shell Systems</u>**
  - **No Problems with Spin Restriction**
- **<u>Sterochemically Flexible</u>**
  - **Large Variety of Structures.**
  - **Little Reorganization**
  - **Facile Ligand Addition/Dissociation**
- **<u>Facilitate Reactions of Bound Ligands</u>**

(Bio)Metals – Perodic Table - Electrons

www.webelements.com

H																	He
Li	Be											B	C	N	O	F	Ne
Na	Mg											Al	Si	P	S	Cl	Ar
K	Ca	**Sc**	***Ti***	**V**	***Cr***	**Mn**	**Fe**	**Co**	**Ni**	**Cu**	**Zn**	Ga	Ge	As	Se	Br	Kr
					Mo												
					W												



Propiedades básicas de un complejo de la proteína metálico

Chem. Rev. 1996, 96, 2239-2314 (1996) RH Holm, P Kennepohl, E I Solomon, Structural and Functional Aspects of Metal Sites in Biology

Prot-L| --------M

δ- δ+

strong attraction



$CH_4 + O_2 + 2e^- + 2H^+ \rightarrow H_3COH + H_2O$

Química en el Centro Catalítico (Sitio activo) de la Enzima de Hierro Metano Monooxigenasa

Ligantes en proteínas – residuos del aminoácido

N



His



Lys

O



Tyr



Glu(+Asp)



Ser

S



Cys



Met

Tetrapirrol está compuesto por 4 unidades de pirrol (porfirinas y ftalocianinas; clorofila, citocromos, pigmentos biliares y vitaminas)



Molybdopterin, un ligante que liga el Mo y el W

JOURNAL of BIOLOGICAL CHEMISTRY (2009) Vol. 284, p. e10, N Kresge, R D Simoni, R L Hill: The Discovery and Characterization of Molybdopterin - the Work of K. V. Rajagopalan





Mo-S-Cu Cluster in CO Dehydrogenase from *Oligotropha carboxidovorans*

$$CO + H_2O \rightarrow CO_2 + 2\,H^+ + 2e^-$$

H Dobbek et al., Proceedings National Academy of Sciences/USA, 99, 15971-15976 (2002)

Geometría – Número de Coordinación

3



Trigonal



Trigonal pyramidal



T-shape

4



Square planar



Tetrahedral

5



Square pyramidal



Trigonal bipyramidal

6



Octahedral

Geometría es importante: Proteínas de Hierro

Tetrahedron | Trigonal Bipyramide | Tetragonal Pyramide | Octahedron





Rubredoxin | 3,4-Protocatechoate Dioxygenase | Tyrosine Hydroxylase | Lipoxygenase

El Solomon, *et al.* (**2000**), Chem. Rev., 100, 235–350

Color y Magnetismo

Alfred Werner (University of Zürich/CH, Nobel Prize in Inorganic Chemistry, 1913)

$Co[(NH_3)_6]Cl_3$ $Co[(NH_3)_5Cl]Cl_2$



Color y Magnetismo

Estados de spin variables de centros metálicos

For a d^5 configuration, Fe(III)



OR



High-Spin, S= 5/2

Low-Spin, S= 1/2

Depending on the METAL ION ENVIRONMENT, balance of Crystal Field Splitting, 10Dq and Spin-Pairing Energy, P



10Dq < P

Weak Field

High Spin



10Dq > P

Strong Field

Low-Spin

Metales – Funciones biológicas

Metal (Ion)	Function, Enzymes
Na	Charge Carrier, Osmolysis/equilibrium
K	Charge Carrier, Osmolysis/equilibrium
Mg	Structure, ATP/ThDP Binding, Photosynthesis,...
Ca	Structure, Signaling, Charge Carrier
V	Nitrogen Fixation, Haloperoxidases, O_2 Carrier
Cr	***Unknown! (glucose metabolism ???)***
Mo	Nitrogen Fixation, Oxidoreductase, O-Transfer
W	Oxidoreductases, Acetylene Hydratase
Mn	Photosynthesis, Oxidases, Structure,...
Fe	Oxidoreductases, O_2 Transport + Activation,e^--Transfer,...
Co	Oxidoreductases, Vitamin B_{12} (Alkyl Group Transfer)
Ni	Hydrogenase, CO Dehydrogenase, Hydrolases, Urease
Cu	Oxidoreductases, O_2 Transport, e^- -Transfer
Zn	Structure, Hydrolases, Acid-Base Catalysis...

Estados de la oxidación de metales en biología

Metal	Valence state (Electron configuration)
Na	Na(I)
K	K(I)
Mg	Mg(II)
Ca	Ca(II)
V	V(V)=(d^0), V(IV)=(d^1), V(III)=(d^2)
Cr	Cr(III)=(d^3),Cr(IV)=(d^2),Cr(V)=(d^1)
Mo	Mo(III)=(d^3),Mo(IV)=(d^2),Mo(V)=(d^1), Mo(VI)=(d^0)
W	W(IV)=(d^2) ,W(V) =(d^1), W(VI)=(d^0)
Mn	Mn(V)=(d^2),Mn(IV) =(d^3),Mn(III) =(d^4),Mn(II) =(d^5)
Fe	Fe(V)=(d^3), Fe(IV)=(d^4),Fe(III)=(d^5),Fe(II)=(d^6), Fe(I)?=(d^7)
Co	Co(III)=(d^6), Co(II)=(d^7), Co(I)=(d^8)
Ni	Ni(III)=(d^7), Ni(II)=(d^8), Ni(I)=(d^9)
Cu	Cu(III)=(d^8), Cu(II)=(d^9), Cu(I)=(d^{10})
Zn	Zn(II) =(d^{10})

Exogenous Ligantes

	Ligand	pK_a
Acid/base	$H_2O/OH^-/O^{2-}$	14,~34
	HCO_3^-/CO_3^{2-}	10.3
	HPO_4^{2-}/PO_4^{3-}	12.7
	H_3CCOO^-/H_3CCOOH	4.7
	HO_2^- /H_2O_2	11.6
	NH_3 /NH_4^+	9.3
	N_3^- /N_3H	4.8
	F^-, Cl^- Br^-, I^-/XH	3.5, -7, -9, -11
Neutral	O_2, CO, NO, RNC	



Modulación de pK_a

$$H_2O + M^{n+} \underset{+H^+}{\overset{-H^+}{\rightleftharpoons}} HO^- \text{-} M^{n+}$$

Metal	pK_a
none	14.0
Ca^{2+}	13.4
Mn^{2+}	11.1
Cu^{2+}	10.7
Zn^{2+}	10.0

4 orders of magnitude !



Control cinético

$$[M^{n+}(H_2O)_m] \underset{+H_2O}{\overset{-H_2O}{\rightleftharpoons}} [M^{n+}(H_2O)_{m-1}]$$

Metal	k (s^{-1})
K^+	1×10^9
Ca^{2+}	3×10^8
Mn^{2+}	2×10^7
Fe^{2+}	4×10^6
Co^{2+}	3×10^6
Ni^{2+}	4×10^4
Fe^{3+}	2×10^2
Co^{3+}	$<10^{-6}$

15 orders of magnitude!

Velocidades de cambio de H_2O

M. Eigen, Nobel Prize Lecture 1967

Expresado como vida de complejos
Útil para mirar la reactividad en
ligand cambian reacciones - catálisis



Estabilidad de complejos del ión metálicos
Irving-Williams Series

H. Irving, R. J. P. Williams (1953) J.Chem.Soc. 3192-3210



Stability order for high-spin divalent metal ion complexes: Maximum at Cu(II), Minimum at Mn(II)

Strong chelating ligand: EDTA



Hexadentate Ligand => strong complexing agent; can be applied to remove metal ions from biological samples (proteins, nucleic acids).

Protein Chelate: Bacterial Metallothionein (MT)



$Zn_4Cys_9His_2$ cluster

C. A. Blindauer et al. (2001) PNAS <u>98</u>, 9593-9598.

- **55 amino acids**
- **One domain**
- **Not only Cys, but also 2 His**
- **Cluster similar to mammalian MT: Essentially a distorted piece of mineral (ZnS)**

Biological Chelate: Siderophores



(a) Enterobactin **(b)** Desferrioxamin

Extremely stable complex of Enterobactin/Fe^{3+} K~ 10^{49}

Release of Fe through a) degradation of ligand, or b) protonation and reduction to Fe^{2+} which binds much weaker to the siderophore.

Modulación de Redox Potentials $E_{1/2}$





$X=O^{-}$: $E_{1/2}$ = -1.21 V

$X=S^{-}$: $E_{1/2}$ = - 0.83 V

R=CH_3 : $E_{1/2}$ = -0.90 V

R=C_2H_5 : $E_{1/2}$ = -0.86 V

R=*i*-Pr : $E_{1/2}$ = -0.74 V

R=*t*-Bu : $E_{1/2}$ = -0.66 V

- **RS^{-} stabilizes Cu(I) state**
- **Positive Potential**

- **El obstáculo de Steric fuerza la geometría tetrahedral, stabilizes Cu(I)**

Metal Site 1: Blue Cu Site (Plastocyanin)

Function: Electron Transfer/Photosynthesis



Covalent Cu-Cys π-bond is mainly responsible for its unique properties

EI Solomon, Inorg. Chem. 2006, 45, 8012-8025



PDB Code: 1PLC
HC Freeman, 1978

Cu(II) Spin-Distribution

Metal site 2: Heme Fe

Function: Electron Transfer/Respiration





Reorganization energy $\propto \Sigma(\Delta R_{ML})^2$

In Cytochromes ≤ 4-5 kcal/mol

Sigfridsson, E.; Olsson, M.H.M.; Ryde, U. (**2001**) *J. Phys. Chem. B*, 105, 5546

Metal Site 3: Iron – Sulfur centers (FeS)

Function: Electron Transfer and Catalysis

Rubredoxin



[3Fe-4S]



[2Fe-2S] Ferredoxin



[4Fe-4S]



[2Fe-2S] Rieske center



Modulación de Redox potentials (H bridges)

2Fe-2S Ferredoxin



$E^{0`}$ ~ -400 mV

2Fe-2S Rieske



$E^{0`}$ ~ +280 mV

(+150 mV without H bridges)

(a) Stephens, P.J.; Jollie, D.R.; Warshel, A. (**1996**) *Chem. Rev.*, 96, 2491

(b) Link, T.A. (**1999**) *Adv. Inorg. Chem.*, 47, 83

Trans-Effect - Modulación de reactividad

A ligand X *trans* to a second ligand Y can influence the stability of the M-Y bond. With X being a strong Lewis base, the M-Y bond will be weakened



Myoglobin
Axial Histidine
O_2 Transport



Cytochrome P450
Axial Cysteine
O_2 Activation

$RH + O_2 + 2H^+ + 2e^- \rightarrow R\text{-}OH + H_2O$

Proteínas modulan las Propiedades de Iones Metálicos

Coordination number

- The lower the higher the Lewis acidity

Coordination geometry

- Proteins can dictate distortion
- Distortion can change reactivity of metal ion

Weak interactions - Second Shell Effects

- Hydrogen bonds to bound ligands
- Hydrophobic residues: dielectric constant can change stability of metal-ligand bonds